

Geometric Quantization

WILLIAM GORDON RITTER*

*Jefferson Physical Laboratory, Harvard University
Cambridge, MA 02138, USA*

February 3, 2008

Abstract

We review the definition of geometric quantization, which begins with defining a mathematical framework for the algebra of observables that holds equally well for classical and quantum mechanics. We then discuss prequantization, and go into details of the general method of quantization with respect to a polarization using densities and half-forms. This has applications to the theory of unitary group representations and coadjoint orbits.

Contents

1	Introduction	2
2	The Mathematical Structure of Physics	2
3	Prequantization	4
4	Quantization	6

*email: ritter@fas.harvard.edu

1 Introduction

The basic problem of quantization is the relationship between observables of classical systems and quantum systems. It is also an opportunity for a bridge to be built between mathematics and physics, since the problem of quantization is motivated by physical concerns, but the technical difficulties involve sophisticated mathematics. Quantum mechanical states are represented by rays in a Hilbert space $\mathcal{H}$, and the observables are represented by symmetric operators on $\mathcal{H}$. In classical mechanics the state space is a symplectic manifold (M, ω) and observables are smooth functions, i.e. elements of $C^\infty(M, \mathbb{R})$.

Taking the view in quantum mechanics that the observables evolve in time while the states remain fixed is known in physics as the Heisenberg picture. The fundamental equation describing the dynamical evolution of a particular (time-dependent) observable A_t is the famous *Heisenberg equation* $\frac{dA_t}{dt} = -\frac{i}{\hbar}[H, A_t]$, where H is the energy observable. This is directly analogous to the situation in classical mechanics. If (M, ω) is the symplectic phase space of a classical system, then the dynamics of a time-evolving observable $f_t : M \times \mathbb{R} \longrightarrow \mathbb{R}$ is given by the differential equation

$$\frac{\partial f_t}{\partial t} = \{H, f_t\}, \quad (1.1)$$

where $\{, \}$ denotes the Poisson bracket. For the canonical choice of symplectic structure on $T^*\mathbb{R}^n$, Eq. (1.1) is equivalent to Hamilton's equations of motion as presented in [1].

The starting point of geometric quantization is to hope that the relationship between Heisenberg's equation and Hamilton's equation exhibited above is a special case of some general situation of deeper mathematical meaning.

2 The Mathematical Structure of Physics

In this section we describe a general mathematical framework for physical theories. Classical mechanics and quantum mechanics are both realizations of this framework; thus, it is an important starting point for quantization. This was inspired in part by lectures of L. Faddeev [2]. The fundamental objects are a set $\mathfrak{A}$ of observables, a set Ω of states, and a *probability interpretation map* $\mathfrak{A} \times \Omega \rightarrow \mathcal{P}$, where $\mathcal{P}$ denotes the set of all nonnegative Lebesgue measurable functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $\int_{-\infty}^{\infty} f(x) dx = 1$ (i.e. probability distributions). For a state η and an observable A , we write the associated probability distribution function as $\eta_A(\lambda)$. Of course, there is a natural mean-value map from $\mathcal{P} \longrightarrow \mathbb{R}$, given by $f \longmapsto \int \lambda f(\lambda) d\lambda$. In all useful examples, $\mathfrak{A}$ and Ω both

have the structure of real vector spaces, and the composition

$$\mathfrak{A} \times \Omega \longrightarrow \mathcal{P} \xrightarrow{\text{mean-value}} \mathbb{R}, \quad \eta, A \longmapsto \langle \eta | A \rangle$$

defines a duality between states and observables. It is clear in physics that certain observables are not independent but rather they are mathematical functions of other, more fundamental observables. An example is the observable p^2 for a classical harmonic oscillator, where p denotes the momentum vector. This fits into the framework above as follows. Given a real function $f : \mathbb{R} \rightarrow \mathbb{R}$ and observables A, B , we write $B = f(A)$ provided that

$$\langle \eta | B \rangle = \int f(\lambda) d\eta_A(\lambda) \quad \text{for all states } \eta$$

In all known cases of practical importance, $\mathfrak{A}$ has the structure of an algebra, and in case $f(x) = \sum \alpha_i x^i$ is a polynomial function, we have $f(A)$, as defined above, equal to $\sum \alpha_i A^i$. Finally, one takes as part of the data a Lie bracket $\{, \}$ on $\mathfrak{A}$ which is an algebra derivation. A fixed observable H , the *Hamiltonian*, is chosen on physical grounds; H equals the total energy and is such that the differential equation $\frac{dA_t}{dt} = \{H, A_t\}$ generates the correct dynamical evolution of observables.

One can reconstruct all features of classical mechanics (even classical statistical mechanics) with the additional assumption that the algebra $\mathfrak{A}$ is commutative. In this situation, there exists a symplectic manifold (M, ω) s.t. $\mathfrak{A} = C^\infty(M)$ as algebras, in which case we take states as normalized measures η on M (a statistical mechanics description), the probability interpretation map as

$$\eta, f \longmapsto \eta_f(\lambda) \stackrel{\text{def}}{=} \int_M \theta(\lambda - f(m)) d\eta(m), \quad \theta = \text{step function},$$

and $\{f, g\} = \omega(X_f, X_g)$ as the Poisson bracket. Nonstatistical mechanics falls out of this by considering a restricted state space (called *pure states*): the space of atomic measures concentrated at a single point of M , which is of course naturally identified with M itself.

In the case of quantum mechanics, the algebra of observables $\mathfrak{A}$ is usually realized as an algebra of linear operators on a complex Hilbert space $\mathcal{H}$, and the space $D(\mathcal{H})$ of positive operators with unit trace (or *density matrices*) is taken as the space of states. In particular, this state space contains the projective Hilbert space (pure states)

$$\mathbb{P}(\mathcal{H}) = \{\text{all projection operators onto 1-dimensional subspaces}\} \cong \mathcal{H} / \sim$$

where $\sim$ is equivalence modulo multiplication by a phase. Elements of $\mathbb{P}(\mathcal{H})$ are known as pure states, while elements of $D(\mathcal{H})$ which cannot be represented as one-dimensional

projectors are known as mixed states. The probability interpretation between a state η and an observable A is given by the pairing

$$\eta_A(\lambda) = \text{Tr}_{\mathcal{H}}(\eta P_A(\lambda))$$

where $P_A(\lambda)$ is the projector function associated to the operator A by the spectral theorem. The dynamical bracket is $\{A, B\} = (i/\hbar)(AB - BA)$, which completes the specification of quantum mechanics in terms of the structure above.

3 Prequantization

Let (M, ω) be a $2n$ -dimensional symplectic manifold and let

$$\epsilon = \left(\frac{1}{2\pi\hbar} \right)^n dp_1 \wedge dp_2 \wedge \cdots \wedge dp_n \wedge dq^1 \wedge dq^2 \wedge \cdots \wedge dq^n$$

be the natural volume element. Based ultimately on physical experiment, Dirac formulated the following prescriptions of the mathematical structure of quantization around 1925, long before mathematicians knew that the procedure was possible. A suitable quantization will produce a quantum mechanical Hilbert space $\mathcal{H}$ from M in a natural way, and will associate to each classical observable $f : M \rightarrow \mathbb{R}$ an operator $\hat{f}$, possibly unbounded, acting on $\mathcal{H}$. On physical grounds, the mapping $C^\infty(M) \rightarrow \mathcal{O}$ given by $f \mapsto \hat{f}$ should at least satisfy the following properties:

(Q1) $f \mapsto \hat{f}$ is $\mathbb{R}$ -linear.

(Q2) if f is constant, i.e. $f(m) = \alpha$ for all $m \in M$ and for some fixed real number α , then $\hat{f} = \alpha I$, where I is the identity operator on $\mathcal{H}$.

(Q3) if $\{f_1, f_2\} = f_3$, then $[\hat{f}_1, \hat{f}_2] = -i\hbar\hat{f}_3$.

If the hat operation is to be a bijective correspondence $C^\infty(M) \xrightarrow{1-1, \text{onto}} \mathcal{O}$, then the Hilbert space needed is too large to be physically meaningful. However, choosing a polarization of M determines a subalgebra of classical observables which can admit bijective quantization maps satisfying Q1-Q3, with the added bonus that the associated Hilbert space is also the space of states of a known quantum mechanical system. We will return to this point.

We also require the *irreducibility postulate*: if $\{f_j\}$ is a complete set of classical observables of (M, Ω) , then the Hilbert space $\mathcal{H}$ has to be irreducible under the action of the set $\{\hat{f}_j\}$. Alternatively, suppose G is a group of symmetries of a physical system both for the classical and quantum descriptions. If G acts transitively on (M, Ω) , then

$\mathcal{H}$ is an irreducible representation space for a $U(1)$ -central extension of the corresponding group of unitary transformations.

Since any symplectic manifold will have a natural volume element ϵ , and hence a natural measure $d\epsilon$, there will also be a natural Hilbert space $\mathcal{H} = L^2(M, d\epsilon)$. Each $f \in C^\infty(M)$ acts on $\mathcal{H}$ by a symmetric operator $-i\hbar X_f$, and this correspondence satisfies Q1 and Q3, but not Q2. However, by modifying this definition appropriately and using a little gauge theory, one arrives at the construction known as *prequantization*, which we describe presently.

Definition 1 *A symplectic manifold (M, ω) is said to be quantizable if ω satisfies the integrality condition, i.e. if the class of $(2\pi\hbar)^{-1}\omega$ in $H^2(M, \mathbb{R})$ lies in the image of $H^2(M, \mathbb{Z})$.*

The integrality condition which appears in Definition 1 is equivalent to the statement that there exists a Hermitian line bundle $B \rightarrow M$ and a connection ∇ on B with curvature $\hbar^{-1}\omega$. It is this latter form of the integrality condition (IC) which we will actually use. In this situation, the space of inequivalent pairs (B, ∇) is parametrized by $H^1(M, S^1)$. This is significant because if M is simply connected, then $H^1(M, S^1)$ is trivial and there is a unique choice of B and ∇ . A bundle $B \rightarrow M$ with connection chosen as above is called a *prequantum bundle*. Let $(\cdot, \cdot)$ be the Hermitian structure on the bundle B , and let $\mathcal{H} = L^2(M, B)$, the space of square-integrable sections of the prequantum bundle, with the inner product $\langle s, s' \rangle = \int_M (s, s') \epsilon$. For $f \in C^\infty(M)$, define a symmetric operator $\hat{f}$ initially on smooth sections of $\mathcal{H}$ by

$$\hat{f}s = -i\hbar \nabla_{X_f} s + f s$$

This choice of $\mathcal{H}$ and of the map $f \mapsto \hat{f}$ satisfies Q1-Q3, but the Hilbert space constructed is too large to represent the space of states of any physically reasonable quantum system. For a function f on M such that the Hamiltonian vector field X_f is complete, the one-parameter group ϕ_f^t of canonical transformations generated by f preserves the scalar product $\langle s, s' \rangle = \int_M (s, s') \epsilon$, and therefore $\hat{f}$ extends to a self-adjoint operator on $\mathcal{H}$. However, if we wish to give a probabilistic interpretation to the scalar product by associating to $\langle \lambda, \lambda \rangle(x)$ the probability density of finding the quantum state described by λ in the classical state described by the point x in classical phase space, we would violate the uncertainty principle since square-integrable sections of B can have arbitrarily small support. Intuitively, a position-space or momentum-space representation corresponds to a certain choice of polarization. Without introducing polarizations, it is no longer true that a wave function sharply peaked in the position variables cannot also be sharply peaked in the momentum variables.

For this reason, the construction outlined above is called *prequantization*, and a refinement of some sort is needed before this procedure can rightly be called “quantization.”

4 Quantization

In quantum mechanics one may represent the Hilbert space as the space of square-integrable complex functions on the spectrum of any complete set of commuting observables. A natural classical analogue of a complete set of commuting observables is a collection of $n = \frac{1}{2} \dim M$ functions $f_1, \dots, f_n$ on M , independent at all points of M where they are defined, such that the Hamiltonian vector fields X_{f_i} are complete, and such that $\{f_i, f_j\} = 0$ for all $1 \leq i, j \leq n$. The Hamiltonian vector fields X_{f_i} span over $\mathbb{C}$ an involutive distribution F such that (i) $\dim_{\mathbb{C}}(F) = \frac{1}{2} \dim(M)$ and (ii) $\omega|_{F \times F} = 0$. A complex distribution F satisfying (i)-(ii) is called a *Lagrangian distribution*. A *polarization* is a complex involutive Lagrangian distribution F such that $\dim(F_x \cap \bar{F}_x)$ is constant over $x \in M$. The complex distributions $F \cap \bar{F}$ and $F + \bar{F}$ are complexifications of certain real distributions traditionally denoted D and E in the literature. The vector spaces given by D and E at a point are ω -perpendicular. The involutivity of F implies that D is involutive, so D is a foliation. We let $\pi_D : M \rightarrow M/D$ denote the projection onto the space of integral manifolds. A polarization F is said to be *admissible* if E is also a foliation, the spaces M/D and M/E are quotient manifolds of M , and the canonical projection $\pi_{DE} : M/D \rightarrow M/E$ is a submersion. For admissible polarizations, the tangent bundle $T\Lambda$ of each integral manifold Λ of D is globally spanned by commuting vector fields. Also, a Hamiltonian vector field X_f lies entirely in D if and only if f is constant along E . Furthermore, each fiber N of π_{DE} has a Kähler structure such that $F|_{\pi_D^{-1}(N)}$ projects onto the distribution of anti-holomorphic vectors on N . For an admissible positive (this means $i\omega(\xi, \bar{\xi}) \geq 0$ for all $\xi \in F$) polarization, the Kähler metric on N is positive definite. A polarization F is said to be *real* if $F = \bar{F}$. Any real-polarized symplectic manifold is locally symplectomorphic to a cotangent bundle with its vertical polarization.

Given a polarization F of (M, ω) with a prequantum line bundle B , one could take sections of B which are covariantly constant along F to form the representation space, except that if λ_1, λ_2 are two such sections, then $\langle \lambda_1, \lambda_2 \rangle$ is constant along D and hence $\int_M \langle \lambda_1, \lambda_2 \rangle \epsilon$ diverges generically unless the leaves of D are compact. Since $\langle \lambda_1, \lambda_2 \rangle$ defines a function on M/D , one could define a scalar product by integrating $\langle \lambda_1, \lambda_2 \rangle$ over M/D , except that there is no canonically defined measure on M/D . The strategy is then to tensor B with another bundle so that $\langle \lambda_1, \lambda_2 \rangle$ may be promoted to a *density* on M/D rather than a scalar function. Tensoring B with $\sqrt{\wedge^n F}$, as we shall see, will

lead to the correct modification of the Bohr-Sommerfeld conditions, and will enable one to construct unitary representations of certain groups of canonical transformations.

The collection of all linear frames of F forms a principal $GL(n, \mathbb{C})$ fibre bundle $\mathcal{B}F$ over M , and associated to this frame bundle is the complex line bundle $\wedge^n F$. Let $ML(n, \mathbb{C}) \xrightarrow{\rho} GL(n, \mathbb{C})$ denote the double covering group of $GL(n, \mathbb{C})$. A *bundle of metilinear frames* of F is a right principal $ML(n, \mathbb{C})$ fibre bundle $\tilde{\mathcal{B}}F$ over M , together with a map $\tau : \tilde{\mathcal{B}}F \rightarrow \mathcal{B}F$ such that the following diagram commutes:

$$\begin{array}{ccc} \tilde{\mathcal{B}}F \times ML(n, \mathbb{C}) & \longrightarrow & \tilde{\mathcal{B}}F \\ \downarrow \tau \times \rho & & \downarrow \tau \\ \mathcal{B}F \times GL(n, \mathbb{C}) & \longrightarrow & \mathcal{B}F \end{array}$$

where the horizontal arrows denote group actions. Let $\chi : ML(n, \mathbb{C}) \rightarrow \mathbb{C}$ denote the unique holomorphic square root of the complex character $\det \circ \rho$ of $ML(n, \mathbb{C})$ such that $\chi(I) = 1$. We define $\sqrt{\wedge^n F}$ to be the fibre bundle associated to $\tilde{\mathcal{B}}F$ with standard fibre $\mathbb{C}$ on which a typical element $C \in ML(n, \mathbb{C})$ acts by multiplication by $\chi(C)$. The space of sections μ of $\wedge^n F$ is isomorphic to the space of complex valued functions on $\mathcal{B}F$ satisfying $\mu^\#(wC) = \det(C^{-1})\mu^\#(w)$ for all $w \in \mathcal{B}_x F$ and $C \in GL(n, \mathbb{C})$, with the isomorphism being $\mu^\# \mapsto \mu(w) \equiv \mu^\#(w_1, \dots, w_n)w_1 \wedge \dots \wedge w_n$. Similarly, the space of sections of $\sqrt{\wedge^n F}$ is isomorphic to the space of functions $\nu^\#$ on $\tilde{\mathcal{B}}F$ satisfying $\nu^\#(\tilde{w}C) = \chi(C^{-1})\nu^\#(\tilde{w})$ for $\tilde{w} \in \tilde{\mathcal{B}}F$ and $C \in ML(n, \mathbb{C})$.

Quantum states of the system under consideration are represented by sections of $B \otimes \sqrt{\wedge^n F}$ which are covariantly constant along F . If σ is such a section, and if ψ is a complex-valued function on M/D which is holomorphic when restricted to fibres of π_{DE} , then $(\psi \circ \pi_D)\sigma$ is also a section of $B \otimes \sqrt{\wedge^n F}$ covariantly constant along F . Thus quantum states may be represented by sections of $B \otimes \sqrt{\wedge^n F}$ which are covariantly constant along D and holomorphic along fibres of π_{DE} .

To each pair (σ_1, σ_2) of sections of $B \otimes \sqrt{\wedge^n F}$ covariantly constant along F , we associate a complex density $\langle \sigma_1, \sigma_2 \rangle_{M/D}$ on M/D . For each $x \in M$, there is a neighborhood $V \ni x$ such that

$$\sigma_i|_V = \lambda_i \otimes \nu_i \quad (i = 1, 2)$$

where λ_i are covariantly constant sections of $B|_V$ and ν_i are covariantly constant sections of $\sqrt{\wedge^n F}|_V$. Consider a basis

$$(v_1, \dots, v_d, u_1, \dots, u_{n-d}, \bar{u}_1, \dots, \bar{u}_{n-d}, w_1, \dots, w_d) \quad (4.1)$$

of $T_x^{\mathbb{C}} M$ such that $\{v_i\}$ is a basis of D_x , $\underline{b} = (v_1, \dots, v_d, u_1, \dots, u_{n-d})$ is a basis of F_x , and for $1 \leq i, j \leq d$ and $1 \leq k, r \leq n-d$ we have

$$\omega(v_i, w_j) = \delta_{ij}, \quad i\omega(u_k, \bar{u}_r) = \delta_{kr} \quad \omega(u_k, w_j) = \omega(w_i, w_j) = 0$$

The basis (4.1) projects under π_D to a basis $\xi_{x,D}$ of $T_{\pi_D(x)}^{\mathbb{C}}M/D$. The value of

$$\langle \lambda_1(x), \lambda_2(x) \rangle \nu_1^{\#}(\tilde{\underline{b}}) \overline{\nu_2^{\#}(\tilde{\underline{b}})} \quad (4.2)$$

(where $\tilde{\underline{b}}$ is a metilinear frame of F at x projecting onto $\underline{b}$) depends only on σ_1, σ_2 and the projected basis $\xi_{x,D}$ of $T_{\pi_D(x)}^{\mathbb{C}}M/D$, hence we define (4.2) to be the value of the density $\langle \sigma_1, \sigma_2 \rangle_{M/D}$ on the basis $\xi_{x,D}$. Hence the sesquilinear form

$$(\sigma_1 \mid \sigma_2)_c := \int_{M/D} \langle \sigma_1, \sigma_2 \rangle_{M/D}$$

is a Hermitian inner product on the Hilbert space $\mathcal{H}^0$ defined as the completion of the pre-Hilbert space of sections σ such that $(\sigma \mid \sigma)_c < \infty$. Note that $\mathcal{H}^0$ is the subspace of the full representation space $\mathcal{H}$ corresponding to the continuous spectrum of the complete set of commuting observables used to define the representation. If the polarization is real and the integral manifolds of D are simply connected, then $\mathcal{H}^0 = \mathcal{H}$.

The complement of $\mathcal{H}^0$ in the representation space $\mathcal{H}$ is spanned by distributional sections of $B \otimes \sqrt{\bigwedge^n F}$ covariantly constant along F . The supports of these sections are restricted by Bohr-Sommerfeld conditions. Let Λ be an integral manifold of D . The operator ∇ of covariant derivative on sections of $B \otimes \sqrt{\bigwedge^n F}$ in the direction F induces a flat connection on $(B \otimes \sqrt{\bigwedge^n F})|_{\Lambda}$. Let $G_{\Lambda} \subset \mathbb{C}^{\times}$ be the holonomy group of this flat connection. The *Bohr-Sommerfeld variety* is

$$S = \{x \in M \mid G_{\Lambda(x)} = 1\}$$

where $\Lambda(x)$ is the integral manifold passing through x . Note that $S = M$ if each Λ is simply connected. Covariantly constant sections of $B \otimes \sqrt{\bigwedge^n F}$ vanish in $M \setminus S$. Thus as claimed the supports of the distributional sections are restricted by Bohr-Sommerfeld conditions. To relate this to the classical Bohr-Sommerfeld conditions, choose a neighborhood U such that $B|_U$ admits a trivializing section λ . Then $\nabla \lambda = -i\hbar^{-1}\theta \otimes \lambda$ where θ is a 1-form on U such that $\omega|_U = d\theta$. For each loop γ in U , the corresponding holonomy is $\exp(i\hbar^{-1} \int_{\gamma} \theta)$. If $\gamma \subset \Lambda$, $\Lambda \in M/D$ then we denote by $\exp(-2\pi i d_{\gamma})$ the element of the holonomy of the flat connection on $(B \otimes \sqrt{\bigwedge^n F})|_{\Lambda}$ corresponding to γ . The condition $G_{\Lambda} = 1$ is then equivalent to

$$\int_{\gamma} \theta = (n_{\gamma} + d_{\gamma})/\hbar, \quad n_{\gamma} \in \mathbb{Z}$$

for each loop γ in Λ .

A polarization F of (M, ω) is said to be *complete* if all Hamiltonian vector fields in F are complete. We will describe the full representation space for a complete admissible real polarization. For $k \in \{0, \dots, n\}$ let

$$M_k = \{x \in M \mid \Lambda_x \cong T^k \times \mathbb{R}^{n-k}\}, \quad \text{and} \quad S_k = S \cap M_k.$$

We note that all integral manifolds of D are isomorphic to products of tori and affine spaces, so $\bigcup_{i=0}^n M_k = M$. For each $x \in S_k$ there exists a neighborhood V of $\pi_D(x)$ in M/D and a codimension k submanifold Q such that

$$\pi_D(M_k) \cap Q \subseteq \pi_D(S_k), \quad \text{and} \quad \pi_D(S) \cap V \subseteq Q \quad (4.3)$$

Let Γ_k denote the space of sections σ of $B \otimes \sqrt{\Lambda^n F}$ satisfying (i) $\text{supp}(\sigma) \subseteq S_k$, (ii) $\pi_D(\text{supp } \sigma) \subset M/D$ is compact, and (iii) for each $x \in S_k$, $\sigma|_{\pi_D^{-1}(Q)}$ is a smooth section of $B \otimes \sqrt{\Lambda^n F}|_{\pi_D^{-1}(Q)}$ covariantly constant along $F|_{\pi_D^{-1}(Q)}$. Let $\sigma_1, \sigma_2 \in \Gamma_k$. By condition (ii), there exist a finite number of disjoint submanifolds $Q_1, \dots, Q_s$ of M/D satisfying (4.3) such that $\text{supp}(\sigma_j) \subseteq \bigcup_{i=1}^s \pi_D^{-1}(Q_i)$. The pair σ_1, σ_2 defines on Q_i a density $\langle \sigma_1, \sigma_2 \rangle_{Q_i}$ by the same procedure as for the continuous part of the spectrum discussed above. The scalar product on Γ_k is defined by

$$(\sigma_1 \mid \sigma_2)_k = \sum_{i=1}^s \int_{Q_i} \langle \sigma_1, \sigma_2 \rangle_{Q_i}$$

and $\mathcal{H}^k$ is defined to be the Hilbert space completion of Γ_k with respect to $(\mid)_k$. The full representation space is a direct sum

$$\mathcal{H} = \bigoplus_{k=0}^n \mathcal{H}^k$$

Quantization of Observables

A quantized operator is said to be *polarized* with respect to a particular polarization P iff it maps polarized sections to other polarized sections. It is necessary to work with polarized operators in order to satisfy the irreducibility postulate. Therefore, we briefly discuss necessary and sufficient conditions for polarized operators. We let $\mathcal{H}_P$ denote a Hilbert space of P -polarized sections coming from a prequantization bundle.

Now $\hat{f}$ maps $\mathcal{H}_P \rightarrow \mathcal{H}_P$ only if the flow of X_f preserves P . In particular, if $\hat{f}\mathcal{H}_P \subset \mathcal{H}_P$ then $[X, X_f] \in V_P(M)$ whenever $X \in V_P(M)$, so only a limited class of observables can be quantized. The elements of $C^\infty(M)$ that can be quantized are precisely the functions of canonical coordinates (p, q) which can be represented locally in the form $f = v^a(q)p_a + u(q)$.

For a given classical observable f , the following conditions are equivalent:

1. $\widehat{f}$ is a polarized operator,
2. $\widehat{f}$ preserves the polarization, in the sense that $[\widehat{f}, \nabla_P]\psi = 0$ for every polarized section ψ , and
3. $[X_f, P] \subset P$, where X_f is the Hamiltonian vector field associated to f .

Cotangent Bundles

In case $M = T^*Q$ with canonical symplectic structure Ω , there is a natural real polarization called the *vertical polarization*, which is spanned by the vector fields $\left\{ \frac{\partial}{\partial p_j} \right\}$. Taking the symplectic potential $\theta = -p_j \partial q^j$, the polarized sections are functions $\psi \in \mathbb{C}(T^*Q)$ such that $\frac{\partial \psi}{\partial p_j} = 0$, i.e. those constant along the fibers of T^*Q , so that $\psi = \psi(q^j)$. The operators corresponding to the observables q_j and p_j are

$$\mathcal{O}_{q^j} = q^j ; \quad \mathcal{O}_{p_j} = -i\hbar \frac{\partial}{\partial q^j}$$

This is known as the *Schrödinger representation* of (T^*Q, Ω) .

Using the polarization spanned by the vector fields $\left\{ \frac{\partial}{\partial q^j} \right\}$ (which is also real), and taking $\theta' = q^j \partial p_j$ as symplectic potential, $\psi = \psi(p_j)$ are the polarized sections and the operators corresponding to q_j and p_j are

$$\mathcal{O}_{q^j} = i\hbar \frac{\partial}{\partial p_j} ; \quad \mathcal{O}_{p_j} = p_j$$

This is the *momentum representation* of (T^*Q, Ω) . The relation between these representations is the Fourier transform.

References

- [1] H. Goldstein, *Classical mechanics*, 2nd ed., Addison-Wesley, Reading, MA (1980).
- [2] P. . Deligne *et al.*, “Quantum Fields And Strings: A Course For Mathematicians. Vol. 1, 2,” *Providence, USA: AMS (1999) 1-1501*.